

ib>-1

ib>-2

國立臺北科技大學九十六學年度碩士班招生考試

系所組別：4210 商業自動化與管理研究所甲組

第一節 統計學 試題

第一頁 共二頁

注意事項：

1. 本試題共九題，配分共 100 分。
2. 請標明大題、子題編號作答，不必抄題。
3. 全部答案均須在答案卷之答案欄內作答，否則不予計分。

1. A random variable X has the following probability density function: $f(x) = \lambda e^{-\lambda(x-a)}$. Please find the moment generating function of X . (5 points)
2. X is a random variable and has distribution $N(\mu, \sigma^2)$. A sample of size 30 from it, say X_1, X_2, \dots, X_{30} , yields the following values: $\sum_{i=1}^{30} X_i = 500$, $\sum_{i=1}^{30} X_i^2 = 8350$. Please find 95-percent (two sided) confidence intervals for μ . (you have to estimate σ first) (5 points)
3. If X_1, X_2, \dots, X_{50} are independent random variables and each of them is distributed as $N(0,1)$. Please find $P[X_1^2 + X_2^2 + \dots + X_{50}^2 > 45]$ (10 points)
4. Suppose that X has a geometric distribution with parameter p . If we let X_1, X_2, \dots, X_n be a random sample from X and let $M = \max(X_1, X_2, \dots, X_n)$ and $K = \min(X_1, X_2, \dots, X_n)$. Please find the probability distribution of M and of K . (10 points)
5. Suppose that $E(Y) = \alpha X^2 + \beta X + \gamma$, where X is pre-defined. Please find the least squares estimates of the parameters of α, β, γ by giving a set of samples $(x_i, Y_i), i=1, 2, \dots, n$. (10 points)
6. Peter uses a computer to do simulation. But this computer, in adding numbers, rounds each number off to the nearest integer. If we know that all rounding errors are independent and uniformly distributed over $(-2, 2)$, please answer the following questions. (15 points)
 - (a) What is the probability that the magnitude of the total error exceeds 30 if 2000 numbers are added?
 - (b) How many numbers may be added together in order that the magnitude of the total error is less than 20, with probability 0.9?

7. In an experiment, a procedure is performed until a particular event A occurs for the first time. On each repetition $P(A)=p$. Suppose that n_1 repetitions are required. Then this procedure is repeated again, and the 2nd time n_2 repetitions are required to yield the event A . If this procedure is conducted for k times, we obtain the sample n_1, n_2, \dots, n_k . Please find the maximum likelihood estimate of p according to the sample. (15 points)
8. Peter is running an investigation. He has a random variable X with distribution $N(\mu, \sigma^2)$. The parameter σ^2 is known. In order to conduct a hypotheses test $H_0: \mu = \mu_0$ vs. $H_1: \mu < \mu_0$, Peter proposed a procedure: obtain a sample of size n and reject H_0 whenever the sample mean $\bar{X} < C$, where C is a constant to be determined. (15 points)
 - (a) Please find an expression for the OC function $L(\mu)$ in terms of normal distribution
 - (b) Please find an expression for C if the significance level of the test is set as $\alpha = 0.01$,
 - (c) Suppose that $\sigma^2 = 4$ and assume that we are testing $H_0: \mu = 30$ vs. $H_1: \mu < 30$. Please find the sample size n and the constant C in order to satisfy the conditions $L(30) = 0.98$, and $L(27) = 0.01$.
9. In typical regression, there are several assumptions that we need to make in order to setup interval estimates and make tests. Please list these assumptions and some possible ways to evaluate them. (15 points)

注意：背面尚有參考資料

TABLE Values of the Standard Normal Distribution Function

$$\Phi(z) = \int_{-\infty}^z \frac{1}{\sqrt{2\pi}} e^{-u^2/2} du = P(Z \leq z)$$

z	0	1	2	3	4	5	6	7	8	9
-3.0	0.0013	0.0010	0.0007	0.0005	0.0003	0.0002	0.0002	0.0001	0.0001	0.0000
-2.9	0.0019	0.0018	0.0017	0.0017	0.0016	0.0016	0.0015	0.0015	0.0014	0.0014
-2.8	0.0026	0.0025	0.0024	0.0023	0.0023	0.0022	0.0021	0.0021	0.0020	0.0019
-2.7	0.0035	0.0034	0.0033	0.0032	0.0031	0.0030	0.0029	0.0028	0.0027	0.0026
-2.6	0.0047	0.0045	0.0044	0.0043	0.0041	0.0040	0.0039	0.0038	0.0037	0.0036
-2.5	0.0062	0.0060	0.0059	0.0057	0.0055	0.0054	0.0052	0.0051	0.0049	0.0048
-2.4	0.0082	0.0080	0.0078	0.0075	0.0073	0.0071	0.0069	0.0068	0.0066	0.0064
-2.3	0.0107	0.0104	0.0102	0.0099	0.0096	0.0094	0.0091	0.0089	0.0087	0.0084
-2.2	0.0139	0.0136	0.0132	0.0129	0.0126	0.0122	0.0119	0.0116	0.0113	0.0110
-2.1	0.0179	0.0174	0.0170	0.0166	0.0162	0.0158	0.0154	0.0150	0.0146	0.0143
-2.0	0.0228	0.0222	0.0217	0.0212	0.0207	0.0202	0.0197	0.0192	0.0188	0.0183
-1.9	0.0287	0.0281	0.0274	0.0268	0.0262	0.0256	0.0250	0.0244	0.0238	0.0233
-1.8	0.0359	0.0352	0.0344	0.0336	0.0329	0.0322	0.0314	0.0307	0.0300	0.0294
-1.7	0.0446	0.0436	0.0427	0.0418	0.0409	0.0401	0.0392	0.0384	0.0375	0.0367
-1.6	0.0548	0.0537	0.0526	0.0516	0.0505	0.0495	0.0485	0.0475	0.0465	0.0455
-1.5	0.0668	0.0655	0.0643	0.0630	0.0618	0.0606	0.0594	0.0582	0.0570	0.0559
-1.4	0.0808	0.0793	0.0778	0.0764	0.0749	0.0735	0.0722	0.0708	0.0694	0.0681
-1.3	0.0968	0.0951	0.0934	0.0918	0.0901	0.0885	0.0869	0.0853	0.0838	0.0823
-1.2	0.1151	0.1131	0.1112	0.1093	0.1075	0.1056	0.1038	0.1020	0.1003	0.0985
-1.1	0.1357	0.1335	0.1314	0.1292	0.1271	0.1251	0.1230	0.1210	0.1190	0.1170
-1.0	0.1587	0.1562	0.1539	0.1515	0.1492	0.1469	0.1446	0.1423	0.1401	0.1379
-0.9	0.1841	0.1814	0.1788	0.1762	0.1736	0.1711	0.1685	0.1660	0.1635	0.1611
-0.8	0.2119	0.2090	0.2061	0.2033	0.2005	0.1977	0.1949	0.1922	0.1894	0.1867
-0.7	0.2420	0.2389	0.2358	0.2327	0.2297	0.2266	0.2236	0.2206	0.2177	0.2148
-0.6	0.2743	0.2709	0.2676	0.2643	0.2611	0.2578	0.2546	0.2514	0.2483	0.2451
-0.5	0.3085	0.3050	0.3015	0.2981	0.2946	0.2912	0.2877	0.2843	0.2810	0.2776
-0.4	0.3446	0.3409	0.3372	0.3336	0.3300	0.3264	0.3228	0.3192	0.3156	0.3121
-0.3	0.3821	0.3783	0.3745	0.3707	0.3669	0.3632	0.3594	0.3557	0.3520	0.3483
-0.2	0.4207	0.4168	0.4129	0.4090	0.4052	0.4013	0.3974	0.3936	0.3897	0.3859
-0.1	0.4602	0.4562	0.4522	0.4483	0.4443	0.4404	0.4364	0.4325	0.4286	0.4247
-0.0	0.5000	0.4960	0.4920	0.4880	0.4840	0.4801	0.4761	0.4721	0.4681	0.4641

TABLE Critical Values for Student's t -Distribution
Pr(Student's $t \leq$ tabled value) = γ

t	0.75	0.90	0.95	0.975	0.99	0.995
1	1.0000	3.0777	6.3138	12.7062	31.8207	63.6574
2	0.8165	1.8856	2.9200	4.3027	6.9646	9.9248
3	0.7649	1.6377	2.3534	3.1824	4.5407	5.8409
4	0.7407	1.5332	2.1318	2.7764	3.7469	4.6041
5	0.7267	1.4759	2.0150	2.5706	3.3649	4.0322
6	0.7176	1.4398	1.9432	2.4469	3.1427	3.7074
7	0.7111	1.4149	1.8946	2.3646	2.9980	3.4995
8	0.7064	1.3968	1.8595	2.3060	2.8965	3.3554
9	0.7027	1.3830	1.8331	2.2622	2.8214	3.2498
10	0.6998	1.3722	1.8125	2.2281	2.7638	3.1693
11	0.6974	1.3634	1.7959	2.2010	2.7181	3.1058
12	0.6955	1.3562	1.7823	2.1788	2.6810	3.0545
13	0.6938	1.3502	1.7709	2.1604	2.6503	3.0123
14	0.6924	1.3450	1.7613	2.1448	2.6245	2.9768
15	0.6912	1.3406	1.7531	2.1315	2.6025	2.9467
16	0.6901	1.3368	1.7459	2.1199	2.5835	2.9208
17	0.6892	1.3334	1.7396	2.1098	2.5669	2.8982
18	0.6884	1.3304	1.7341	2.1009	2.5524	2.8784
19	0.6876	1.3277	1.7291	2.0930	2.5395	2.8609
20	0.6870	1.3253	1.7247	2.0860	2.5280	2.8453
21	0.6864	1.3232	1.7207	2.0796	2.5177	2.8314
22	0.6858	1.3212	1.7171	2.0739	2.5083	2.8188
23	0.6853	1.3195	1.7139	2.0687	2.4999	2.8073
24	0.6848	1.3178	1.7109	2.0639	2.4922	2.7969
25	0.6844	1.3163	1.7081	2.0595	2.4851	2.7874
26	0.6840	1.3150	1.7056	2.0555	2.4786	2.7787
27	0.6837	1.3137	1.7033	2.0518	2.4727	2.7707
28	0.6834	1.3125	1.7011	2.0484	2.4671	2.7633
29	0.6830	1.3114	1.6991	2.0452	2.4620	2.7564
30	0.6828	1.3104	1.6973	2.0423	2.4573	2.7500
31	0.6825	1.3095	1.6955	2.0395	2.4528	2.7440
32	0.6822	1.3086	1.6939	2.0369	2.4487	2.7385
33	0.6820	1.3077	1.6924	2.0345	2.4448	2.7333
34	0.6818	1.3070	1.6909	2.0322	2.4411	2.7284
35	0.6816	1.3062	1.6896	2.0301	2.4377	2.7238
36	0.6814	1.3055	1.6883	2.0281	2.4345	2.7195
37	0.6812	1.3049	1.6871	2.0262	2.4314	2.7154
38	0.6810	1.3042	1.6860	2.0244	2.4286	2.7116
39	0.6808	1.3036	1.6849	2.0227	2.4258	2.7079
40	0.6807	1.3031	1.6839	2.0211	2.4233	2.7045
41	0.6805	1.3025	1.6829	2.0195	2.4208	2.7012
42	0.6804	1.3020	1.6820	2.0181	2.4185	2.6981
43	0.6802	1.3016	1.6811	2.0167	2.4163	2.6951
44	0.6801	1.3011	1.6802	2.0154	2.4141	2.6923
45	0.6800	1.3006	1.6794	2.0141	2.4121	2.6896